Nozomi F1

Cabbage







Nozomi F1

Segment

Maturity

Harvest season

Head weight

Head colour

Resistances

Features

Early

Spring

1.0-1.5 kg

Light green

spring harvests

Blackrot

55-65 Days after

transplanting (DAT)

Intermediate Resistance to

Dense uniform heads for

Brassica oleracea L.

the second s	
AN A STATE OF A STATE	
Contraction of the second s	
State and Ba	
and the second	
and the second	
and the second	
Carlos and the second second second	
A CONTRACT AND A CONTRACT OF A CONTRACT OF	
A STATE AND A STATE AN	
AND IN COMPLETE AND ADDRESS OF THE OWNER.	
AND AND ADDRESS OF ADDRESS OF ADDRESS OF ADDRESS ADDRESS ADDRESS ADDRESS ADDRESS ADDRESS ADDRESS ADDRESS ADDRES	
and the second	
AND REPORTS ADDRESS AND REPORTS	
AND INCOMENTS OF A DESCRIPTION OF A DESC	
A REAL PROPERTY AND A REAL PROPERTY A	
and the second se	
and the second	
and a supervision of the supervi	
A STATE OF A	

Closest Sakata variety	- Candisa - Charmant
Advantages	 Excellent field holding ability and resistance to cracking Uniform size, high rate of marketable heads
Advices	- For cold to warm

DISCLAIMER: This information is based on our observations and/or information from other sources. As crop performance depends on the interaction between the genetic potential of the seed, its physiological characteristics, and the environment, including management, we give no warranty express or implied, for the performance of crops relative to the information given nor do we accept any liability for any loss, direct or consequential, that may arise from whatsoever cause.



